

# PHYSICAL THERAPY SECTION WILLIAM BEAUMONT ARMY MEDICAL CENTER 5005 N. PIEDRAS ST EL PASO, TEXAS 79920



# TOTAL HIP ARTHROPLASTY REVISION, LATERAL APPROACH

Revised August 2008

#### **Discharge Criteria**

### To Rehabilitation Hospital

Independent/minimum assist with bed mobility Ambulation with assistive device for 30 feet

#### To Home

Independent bed mobility and transfers
Able to safely negotiate home obstacles such as stairs and carpet
Independent ambulation with assistive device for 300 feet
Coordinate with Social Work Services to obtain home equipment:

— Wheeled walker, Three-in-one, Reacher

#### **Precautions x 6 Weeks**

No Hip Flexion > 60 degrees
Wear TED Hose
Sleep on back
Pillow under ankle, NOT under knee, keep foot of bed flat
Pillow between legs while sleeping
No active Abduction exercises
No straight leg raise (SLR)
No ER > 30 degrees
No Extension > 30 degrees
No Adduction past midline

#### Weeks 1 - 6

Walker or crutches – flat foot weight bearing (FFWB) with up to 20# pressure Ankle pumping
Heel slides, AROM, AAROM with above restrictions
Quad sets, Co-contractions quads/hams
Short arc quads with up to 10#

#### Goals

Protection Hip ROM 0-60 degrees

#### Weeks 6 - 12

Walker or crutches – partial weight bearing (PWB) up to 50%

No hip flexion > 90 degrees

Continue to sleep on back

Continue pillow between legs while sleeping

Continue appropriate previous exercises

Progressive abductor strengthening

- In standing
- Sidelying
- Add light weight in sidelying when able to perform 25 reps

Hip extension - Standing or prone

Sitting knee extension (chair or mat) 90-0 degrees

Weight shifts in parallel bars

Mini squats 0-45 degrees in parallel bars

Double leg heel raises

Goal - Hip ROM 0-90 degrees

#### **Months 3 - 4**

Walker or crutches – weight bearing as tolerated (WBAT)

- Progress to cane and D/C when gait is WNL

Hip flexion > 90 degrees

Continue appropriate previous exercises

Progressive abductor strengthening

- In standing with Theraband, bilaterally
- Hip weight machine x 4 bilaterally

Forward and retro walking in parallel bars

Forward, retro and lateral step downs (small step)

Wall squats

Straight leg raise (SLR)

Leg press (< 90 degrees hip flexion) up to ½ body weight

Hamstring curl weight machine

Knee extension weight machine

Single leg (stork) standing

Single leg heel raises

Stationary bicycle

Stretches - Hamstring, quads, hip flexors, ITB

#### Goals

Symmetrical hip ROM

Normal gait

#### Months 4-5

Continue appropriate previous exercises with increasing resistance as tolerated Practice sit-to-stand without using hands

Stair training

Proprioception exercises as tolerated (age dependent)

Continued on following page

Treadmill – Walking progression program

Elliptical trainer Pool therapy

#### Goals

Walk x 20 minutes Stand from sitting without use of hands

## <u>Months 5 - 6</u>

Discontinue supervised PT Resume all recreational activities as tolerated Encourage non-impact activities See attached Precautions and Activities

BRYAN L. BOYEA LTC, SP Chief, Physical Therapy MICHAEL S. MACHEN LTC(P), MC Chief, Orthopedics